

super fruits

blueberries, raspberries, blackberries, strawberries

▶ Are you looking for enhanced robustness, higher flowering rates and higher yield?

▶ Would you like to achieve bigger, more uniform in size and shape fruits, with full flavor and aroma?

Apply

Amino16

Amino16 B Zn

Amino16 Si Zn

by
EVYP



Amino16®



Applications:

1. Beginning of the emergence of the young shoots.
2. Right before the first flowering.
3. Right after fruit setting of the 1st flowering stage.
4. At the changing of the color, before the 1st harvesting.
5. After harvesting and before the leaves fall. (The leaves should still be green).



Dosages: 10 lt per Ha through the irrigation network.
Alternatively, 5 lt per 1000 lt of water with foliar spraying



www.origami.com.gr

Amino 16® is suggested for the 1st, 3rd and 4th application.

Amino 16® with Boron (B) and Zinc (Zn) is suggested for the 2nd and 5th application. **Amino 16® with Silicon (Si) and Zinc (Zn)** can be used after high temperature fluctuations or intense rainfalls.

Amino16® B Zn



Amino16® Si Zn



For more information about each cultivation, please visit our website: www.evyp.gr

EVYP

A5 Road
Industrial Area of Thessaloniki
PC 57022, Greece
T/F: +30 2310 795 285
info@evyp.gr / www.evyp.gr

